INDEX TO VOLUME V.

ACANTHIS brewsterii, 401. exilipes, 413. hornemannii, 413. linaria, 358. Accentor, Golden-crowned, 315. Accipiter atricapillus striatulus, 145. cooperi, 20, 145, 246, 271, 316, velox, 20, 145, 271, 350. Actitis macularia, 20, 145, 149, 151, 270, 349, 389, 433. Actodromas couesi, 197. Æchmophorus clarkii, 401. occidentalis, 15, 140, 201. Ægialitis meloda circumcincta, 147. mongola, 197. nivosa, 110. pyrrhothorax, 197. semipalmata, 327. vocifera, 20, 145, 270, 349. wilsonia, 186. Æstrelata hasitata, 81. jamaicensis, 81. Agelaius gubernator, 139. phœniceus, 22, 150, 152, 154,

207, 262, 272, 315, 389.

phæniceus sonoriensis, 397.

phœniceus bryanti, 397.

tricolor, 262.

Aimophila cahooni, 93, 94. mcleodii, 92, 94. rufescens, 94. Aix sponsa, 18, 65, 142, 269, 346,

Wandering, 394.

Yellow-nosed, 107, 318.

torda, 134, 135, 281, 382.

Agyrtria alleni, 263.

378, 389. Ajaja ajaja, 269.

Alauda arvensis, 180. Albatross, Spectacled, 394.

Alca impennis, 329.

subspecies of the Seaside Sparrow (Ammodramus maritimus), 284; description of a new species of the genus Tityra from Ecuador, 287; notes on Louisiana birds, 324; notice of his paper 'On the Structure of Birds in relation to Flight,' 418; note on the correct name of Symphemia semipalmata inornata Brewst., 423; further notes on Seaside Sparrows, 426; see also Faxon, W., and Scott, W. E. D. Amaurospiza æquatorialis, 411. axillaris, 412. American Ornithologists' Union, fifth meeting of the, 95. Ammodramus australis, 274, 398. caudacutus, 422. caudacutus nelsoni, 422. caudacutus subvirgatus, 397. henslowii, 22, 210, 389, 390. leconteii. 210. maritimus, 285, 286, 426. maritimus peninsulæ, 287, 398, 426. maritimus sennetti, 286, 287, 398, 426. nigrescens, 284, 285. sandwichensis, 358. sandwichensis alaudinus, 358. sandwichensis brunnescens, 265. sandwichensis savanna, 150, 152, 154, 273, 326. savannarum passerinus, 22, 273. savannarum perpallidus, 15. Ampelis cedrorum, 23, 31, 150, 152,

275, 360, 389. Anas acuta, 62.

Allen, F. H., Tameness of the Pine

Allen, J. A., descriptions of two new

Siskin, 426.

Anas albeola, 67. americana, 63, 141, 269, 346, 378. arborea, 59. autumnalis, 60. bahamensis, 62. boschas, 17, 61, 141, 268, 312, 313, 314, 316, 346, 378. cæ rulescens, 59. canadensis, 59. carolinensis, 17, 64, 141, 269, 346, 378. clypeata, 64. collaris, 66. cyanoptera, 141. discors, 17, 63, 269, 346, 378. dominica, 68. fimbriata, 62. (Fuligula) rufitorques, 66. fulvigula, 61, 108, 378, 444. ilathera, 62. jacquini, 59. marila, 65. maxima, 61. mexicana, 64. obscura, 61, 268, 312, 346, 389, 390, 444. perspicillata, 67 (Querquedula) discors, 64. rubida, 68. rubirostris, 62. spinosa, 68. sponsa, 65. strepera, 61, 141, 173, 268. urophasianus, 62. vallisneria, 66. viduata, 60. Anhinga anhinga, 71, 268, 377. Anhinga, 268. 377 Anous stolidus, 80. Anser albifrons, 58, 311. albifrons gambeli, 58,142,143, 311. cærulescens, 59. gambeli, 58. hyperboreus, 58. pictus, 311. Anthus cervinus, 307, 410. pensilvanicus, 159, 276, 362, 434pratensis, 307. spraguei, 15 Antrostomus carolinensis, 272. vociferus, 21, 149, 151, 272, 389. Aphalus capensis, 292, 293, 301. Aphelocoma californica hypoleuca, 397. insularis, 396. Aquila chrysaëtos, 145, 313, 315, 350.

Aramides maculatus, 52. Aramus giganteus, 50, 51. guarauna, 50. pictus, 51. scolopaceus, 51. scolopaceus giganteus 51. Archibuteo lagopus sancti-johannis, 145, 350. Ardea bahamensis, 83. cærulea, 268, 269. candidissima, 268,269. egretta, 19, 143, 268, 269, 348. exilis, 50. herodias, 143, 183, 268, 269, 347· lentiginosa, 49. nævia, 49. nycticorax, 49. occidentalis, 195. pealei, 184, 401. purpurea, 196. rufa, 184. rufescens, 184. tricolor ruficollis, 268, 269. violacea, 48. virescens, 19, 84, 144, 269, 348. virescens frazari, 83, 394. wardi, 183, 195. wuerdemanni, 195, 394. Ardeola exilis, 50. Ardetta eurhythma, 196. exilis, 50. neoxena, 394. Arenaria interpres, 110, 147. Asio accipitrinus, 146, 350. wilsonianus, 20, 146, 350. Atkinson, G. F., notice of his 'Preliminary catalogue of the birds of North Carolina, 311. Auk, Great, 278, 330, 331, 380, 422. Auklet, Whiskered, 410. Auriparus flaviceps, 166. Averill, C. K., Jr., feeding habits of Sitta canadensis, 118; the Yellow Rail in Connecticut, 319. Avocet, 16 Aythya affinis, 65, 142, 203, 269, 346, 379. americana, 66, 142. collaris, 66, 142, 269, 379. marila nearctica, 18, 142, 379. vallisneria, 66, 142. Aythyia vallisneria. 66. BAIRD, Spencer Fullerton, morial address upon, 1. Baldpate, 269, 346, 378. Bartramia longicauda, 145, 270, 389.

Basileuterus culicivorus, 450. culicivorus godmani, 450. godmani, 450.

Becard, Xantus's, 425.

Beckham, C. W., occurrence of the Florida Blue Jay (Cyanocitta cristata florincola) in Southwestern Texas, 112; a Philadelphia Vireo and a cobweb, 115; death of, 445.

Bendire, C. E., eggs of the Ivory Gull (Gavia alba), 202; notes on the habits, nests, and eggs of the genus Sphyrapicus Baird, 225; notes on the nest and eggs of Peucæa æstivalis bachmani Aud., Bachman's Sparrow, 351; notes on the habits, nests and eggs of the genus Glaucidium Boie, 366.

Bennett, A. L., Surf Scoter (Oidemia perspicillata) in Kansas,

203.

Berlepsch, H. v., descriptions of new species and subspecies of birds from the neotropical region, 449.

Berlepschia, 194.

Bernicla canadensis, 59.

Bird, Inca, 27. Man-of-war, 170, 173, 319, 378. Red-billed Tropic, 241.

Bittern, 268.

American, 269, 347. Least, 19.

Blacicus flaviventris, 158. martinicensis, 158.

Blackbird, Brewer's, 15.

Crow, 15, 113. Red-winged, 22, 150, 272, 315. Rusty, 273.

Yellow-headed, 362.

Blue Peter, 270.

Bluebird, 22, 25, 151, 153, 277. Mountain, 168.

Western, 167. Bobolink, 150, 151, 272.

Bob-white, 20, 317, 349. Cuban, 395.

Florida, 270. Bogdanow, M. N., death of, 333. Bonasa umbellus, 20, 149, 151, 153,

314, 338, 339, 340, 349, 389. umbellus sabini, 145.

Booby, Blue-footed, 241. Brewster's, 241, 242.

Botaurus exilis, 19.

lentiginosus, 49, 143, 268, 269, 328, 347.

Botaurus minor, 49. Brachyrhamphus kittlitzii, 410.

Brant, Black, 143.

Branta canadensis, 17, 59, 143, 347. canadensis hutchinsii, 143. canadensis minima, 143. canadensis occidentalis, 143. hutchinsii, 311.

nigricans, 143.

Brewster, W., descriptions of sup-posed new birds from Lower California, Sonora, and Chihuahua, Mexico, and the Bahamas, 82; the Prairie Horned Lark (Otocoris alpestris praticola) on the coast of Massachusetts, III; on three apparently new subspecies of Mexican birds, 136; Quiscalus quiscula aglæus at Charleston, South Carolina, 208; breeding of the Golden-crested Kinglet (Regulus satrapa) in Worcester County, Massachusetts, with a description of its nest and eggs, 337; notes on the birds of Winchendon, Worcester County, Massachusetts, 386; a second Maine (and fourth New England) specimen of Swainson's Hawk (Buteo swainsoni), 424; Cistothorus pal-ustris marianæ on the coast of Georgia, 432; see also Merrill, I. C.

Brodie, W., bird notes from Toron-

to, 211.

Brown, H., *Ionornis martinica* in Arizona, 109; on the nesting of Palmer's Thrasher, 116.

Brown, J. C., Carpodacus purpur-eus at Portland, Maine, in winter, 200.

Bubo virginianus, 18, 271, 313, 351,

389. virginianus subarcticus, 146.

Bucephala albeola, 67.

Buffle-head, 347, 379. Bullfinch, Cassin's, 410.

Bunting, Bay-winged, 150.

Beautiful, 399.

Cow, 207. Indigo, 22, 152, 274.

Bush-tit, Lead-colored, 166. Lloyd's, 43, 400. Butalis griseisticta, 197.

Buteo borealis, 18, 149, 223, 271,

350, 389.

borealis calurus, 145, 205. cooperi, 401.

Buteo fuliginosus, 185, 401. latissimus, 20, 350. lineatus, 20, 271, 350. pennsylvanicus, 247. swainsoni, 145, 424.

Butler, A. W., on a new subspecies of *Ammodramus sandwichensis* from Mexico, 264.

Buzzard, Turkey, 245, 248, 316, 350. CAIRINA moschata, 63.

Calamospiza melanocorys, 438-444. Calcarius lapponicus, 181, 438, 440,

ornatus, 14, 321. Calidris arenaria, 179.

Callipepla elegans bensoni, 194. Calliste cucullata, 157.

Campephilus principalis, 185, 186. Campylorhynchus brunneicapillus, 116, 162.

zonatus, 449. zonatus costaricensis, 449.

Carvasback, 142. Carbo mexicanus, 71. Cardellina rubrifrons, 36, 385.

Cardinal, 22, 119, 274. Texas, 119.

Cardinalis cardinalis, 22, 119, 274, 421.

Carduelis elegans, 211. magellanicus, 412.

Carpodacus cassini, 412.
frontalis, 412.
frontalis rhodocolpus, 401.
frontalis ruberrimus,397.
hæmorrhous, 412.
mexicanus frontalis, 401.
purpureus, 12, 150, 152, 154,

209, 273, 389. purpureus californicus, 357. roseipectus, 412.

Carpodectes antoniæ, 194. Cassique, Yellow-tailed, 27. Catbird, 24, 151, 153, 277, 388.

Catharista atrata, 270. Cathartes aura, 18, 145, 179, 245, 248, 270, 316, 350.

Catherpes mexicanus conspersus, 162.

Cedarbird, 150, 152. Centrocercus urophasianus, 367. Centurus blakei, 159.

caymanensis, 159. nyeanus, 159.

Ceophlœus pileatus, 111, 149, 240, 271, 313, 389, 390.

Cepphus mandtii, 311 motzfeldi, 401.

Certhia familiaris americana, 19,

Certhia familiaris mexicana, 165. familaris occidentalis, 362. mexicana, 450. mexicana albescens, 450.

Certhiola sharpei, 157.

Ceryle alcyon, 21, 149, 151, 153, 251, 271.

Chadbourne, A. P., Puffinus borealis at Gardiner's Bay, N. Y., 202; Molothrus ater again in Massachusetts in winter, 207.

Chætura brachyura, 158. cinereiventris, 158. pelagica, 21, 149, 151, 154, 272, 389. vauxi, 256.

Chamberlain, M., the Yellow-nosed Albatross (Thalassogeron culminatus) in the Gulf of St. Lawrence, 107; notice of his 'Catalogue of Canadian birds,' 189; notice of his 'Systematic Table of Cana-

dian Birds, 198.
Chapman, F. M., observations on the nocturnal migration of birds, 37; a list of birds observed at Gainesville, Florida. 267; Oidemia perspicillata in Florida, 319; birds at Aiken, S. C., 324; list of additions to the North American avifauna and of eliminations and changes in nomenclature proposed since the publication of the A. O. U. Check-List, 393.

Charadrius dominicus, 308-310. dominicus fulvus, 309, 310. fulvus americanus, 192. fulvus dominicus, 192. fulvus virginicus, 192. squatarola, 110. virginicus, 309.

Charitonetta albeola, 67, 142, 203, 347, 379.

Chasiempsis dolei, 197. Chat, Long-tailed, 36.

Yellow-breasted, 24, 116, 405. Chaulelasmus streperus, 61. Chelidon erythrogaster, 23, 31,

150, 152, 154, 360, 389. Chen cærulescens, 59, 311, 394. hyperborea, 58, 142.

hyperborea, 58, 142. hyperborea nivalis, 183, 379. Chickadee, 151, 153.

Ckadee, 151, 153.
Carolina, 25, 277.
Hudsonian, 410.
Kowak, 400.
Mountain, 118.
Plumbeous, 46, 400.
Siberian, 410.

Chicken, Prairie, 349. Chlæphaga poliocephala, 198. Chondestes grammacus, 186, 438. Chordeiles popetue, 37, 154.

popetue chapmani, 37, 396. henryi, 37. sennetti, 37, 396. virginianus, 21, 149, 151, 256,

314, 389. virginianus chapmani, 186, 272, 396. virginianus henryi, 256.

virginianus sennetti, 256, 396. Chræcocephalus atricilla, 75. Chroicocephalus atricilla, 75. Chrysomitris boliviana, 412.

capitalis, 412. icterica, 412. longirostris, 412. magellanica, 412. notata, 412. sclateri, 412.

siemiradzkii, 412. stejnegeri, 412. xanthogastra, 412.

Chrysotis caymanensis, 159. Chuck-will's-widow, 272. Cinclus mexicanus, 362.

Circus hudsonius, 145, 151, 271, 350.

Cistothorus marianæ, 188, 400. palustris, 24. palustris marianæ, 432. palustris paludicola, 165, 362,

400.
Clangula albeola, 67.
americana, 67.
glaucion, 67.
glaucion americana, 67.
glaucium, 67.
hvemalis, 203.

Clark, H. L., notice of his 'Birds of Amherst.' 105.

Clivicola riparia, 23, 31, 152, 154, 329, 389.

Coccothraustes vespertina, 113, 114, 208, 357, 425, 439, vulgaris, 439.

Coccyzus americanus. 20. 271. americanus occidentalis, 396. erythrophthalmus, 21, 149, 151, 389. maynardi, 159, 396.

minor, 185. Colaptes auratus, 21, 149, 151, 154, 271, 325, 389. cafer, 16, 240, 256. gundlachi, 159. Coleman, G. A., Coccothraustes vespertina in Nebraska, 425. Colinus graysoni, 401.

ridgwayi, 401. virginianus, 20, 313, 316, 349. virginianus cubanensis, 395. virginianus floridanus, 270.

Collocalia fuciphaga, 333, 335. Columba fasciata vioscæ, 86, 395. leucocephala, 185.

Columbigallina passerina, 270. passerina pallescens, 395.

Colymbus auritus, 373. carolinensis, 156. dominicensis, 155. dominicus, 155. holboellii. 156. podiceps, 156.

podiceps, 156.
Comeau, N. A., the Yellow-nosed
Albatross (Thalassogeron culminatus), 318.

Compsothlypis americana. 24, 150, 152, 275, 326, 389, 431, 433. Contopus borealis, 148, 151, 154,

257, 389, 390. richardsoni, 257, 258. virens, 21, 150, 151, 154, 272, 389.

Conurus carolinensis, 148, 185. Coot, 267, 273,

American, 270, 348. Corethrura cayennensis, 54. guatemalensis, 54.

Cormorant, Florida, 377. Corvus americanus, 21. 150, 151,

154, 179, 312, 316, 389, 414, americanus floridanus, 272. americanus hesperis, 262, 397. corax carnivorus, 397. corax principalis, 397.

corax sinuatus, 198, 241, 262, 329. corone orientalis, 196.

levaillantii, 196. ossifragus, 21, 272.

Cory, C. B., description of a supposed new form of *Margarops* from Dominica, 47; an apparently new *Elainea* from Barbadoes, West Indies, 47; the birds of the West Indies, including the Bahama Islands, the Greater and the Lesser Antilles, excepting the Islands of Tobago and Trinidad, 48, 155; the European Kestrel in Massachusetts—a correction, 205; descripters.

tion of a new Myiarchus from the West Indies, 266.

Corydomorphæ, 207. Cotinga ridgwayi, 194. Coturnix coturnix 314.

Coues, E., new forms of North American *Chordiles*, 37; notice of his 'Key to North American Birds,' 101; note on Rostratul-inæ, 204; Corydomorphæ, 207; notes on the nomenclature of the muscles of volation in birds'

wings, 435. Coues, E. B., nesting of the Prairie Warbler (Dendroica discolor) in the vicinity of Washington, D. C., 405.

Cowbird, 15, 21, 151, 272.

Crane, Sandhill, 205, 348. Creciscus jamaicensis, 55.

Creeper, Black-and-white, 150, 433. Brown, 19, 153.

Mexican, 165.

Crex crex, 177. galeata, 56.

minuta, 54. Crithagra brasiliensis, 412. Crœcocephalus minutus, 171.

Crossbill, 324, 325. American, 22, 152, 317.

Mexican, 114.

Red, 115, 208, 340. White-winged, 152, 410. Crow, 150, 179, 312, 316.

American, 21, 151. California, 397. Clarke's, 262. Fish, 21, 272.

Florida, 272. Crybastus gossii, 54.

minutus, 55 Cuckoo, Black-billed, 21, 149, 151.

California, 396. Mangrove, 185. Maynard's, 396.

Yellow-billed, 20, 271. Cupidonia cupido brewsteri, 402.

Curlew, 14.

Cyanocephalus cyanocephalus, 262, 413.

Cyanocitta cristata, 21, 150, 151. 154, 389, 434.

cristata florincola, 112, 272, 324.

stelleri, 261. stelleri annectens, 396. stelleri frontalis, 261.

Cyanolesbia nehrkorni, 200.

Cyanopterus clypeata, 65. discors, 63. inornatus, 63.

Cyanospiza cyanella, 412. Cypseloides niger, 424.

Cypselus apus, 329. DAFILA acuta, 62, 142, 269, 316, 346, 378. bahamensis, 62.

urophasianus, 62. Daption capensis, 198.

Davison, J. L., breeding of the Cerulean Warbler (Dendroica cærulea) in Niagara County, New York, 430.

Demiegretta ringeri, 196.

Dendragapus canadensis, 317. obscurus fuliginosus, 145, 367. Dendrocincla castanoptera, 195.

lafresnayei, 195. rufo-olivacea, 194.

Dendrocygna arborea, 59. autumnalis, 60. viduata, 60.

Dendrocygnus arborea, 60. spinosa, 68. sponsa, 65. viduata, 60.

Dendroica æstiva, 16, 21, 24, 150, 152, 154, 389, 433.

æstiva morcomi, 34, 138, 361, 399. æstiva sonorana, 137, 400.

auduboni, 16, 34, 361. blackburniæ. 150, 153, 326, 389, 391, 433.

cærulea, 323, 325, 430, 431. cærulescens, 149, 150, 153, 154, 276, 326, 389, 391, 433. carbonata, 401.

castanea, 153, 326, 433.

coronata, 16, 34, 149, 150, 153, 154, 276, 326, 388, 389, 391, 430, 433. discolor, 22, 24, 276, 405.

dominica, 276. dominica albilora, 276.

graciæ, 34. kirtlandi, 148.

maculosa, 153, 326, 388, 389, 391, 433-

nigrescens, 34, 104, 361. occidentalis, 35, 361. olivacea, 34.

palmarum, 170, 182, 276, 324, 326, 433.

palmarum hypochrysea, 182, 276.

Dendroica pennsylvanica, 150, 153, 154, 326, 389, 433. striata, 153, 182, 276, 326, 432. tigrina, 187, 211, 275. townsendi, 35, 361. vigorsii, 24, 276, 326, virens, 150, 153, 154, 326, 389, 391, 433.

vitellina, 157.
Dickcissel, 23.
Die Dipper, 268.
Diòmedea exulans, 394.
melanophrys, 394.
Diplopteryx chirurgus, 292.
Diver, Red-throated, 171.
Dolichonyx oryzivorus, 150, 151,

154, 272, 314, 389.
Dove, 270.
Carolina, 119.
Ground, 270.
Mexican Ground, 395.
Mourning, 20, 270, 349.

Turtle, 17.

Dowitcher, 110.

Long-billed, 147, 184.

Dryobates borealis, 271.

pubescens, 16, 21, 151, 154,

271. pubescens gairdnerii, 240, 251, 252. villosus, 21, 149, 151, 154, 389, 390.

390. villosus audubonii, 271. villosus harrisii, 213, 214, 218, 240, 251. villosus hyloscopus, 252.

Duck, American Scaup, 18.

Black, 108, 268, 346, 390, 444.

Blue Mountain, 81.

Bufflehead, 203.

Florida, 378, 444.

Florida Black, 268.

Gadwall, 173.

Golden-eye, 203.

Isle of Shoal, 175.

King Eider, 175.

Lesser Black-head, 203.

Lesser Scaup, 269, 346, 379.

Long-tailed, 203.

Ring-necked, 269, 379.

Ruddy, 269, 347, 379, 416.

Shoveller, 17.

Wood, 18, 269, 346, 379.

Dutcher, W., bird notes from Long Island, N. Y., 169. *

Dwight, J., Jr., two records for Long Island, New York, 324.

Dysporus cyanops, 71.

Dysporus fiber, 72. hernandezi, 73. leucogaster, 72. piscator, 73. sula, 72. EAGLE, Bald, 18, 149, 271, 350.

Golden, 313, 315, 350.

Eames, E. H., notes on *Helmin-thophila leucobronchialis*, 427;

Dendroica cærulea at Seymour,

Connecticut, 431.

Ectopistes migratorius, 270, 314, 349, 389. Egret, American, 19, 269, 348.

Egret, American, 19, 269, 34 Great White, 268. Little White, 268. Reddish, 184.

Eider, King, 174. Elainea barbadensis, 47, 158. pagana, 158.

Elanoides forficatus, 270. Elliot, D. G., description of a new species of Hummingbird of the genus Agyrtria, 263; the Jacanidæ, 288.

Emberiza aldunatei, 194. cyanella, 412. Embernagra rufivirgata, 352, 442. Emmet, R. T., the Black Duck in Chihuahua, 108.

Empidonax acadicus, 21, 272, 422. cineritius, 90, 396. difficilis, 90, 91, 258. flaviventris, 3, 151. hammondi, 258. minimus, 3, 150, 151, 389, 434. obscurus, 258. pusillus, 258. pusillus traillii, 151, 154, 422.

Eremophila alpestris, 111.
Eremotes occidentalis. 184.
pusillus, 185. 349.

Erismatura dominica, 68. rubida, 68, 142, 269, 347, 379, 416. spinosa, 68.

Erythra minuta, 54.
Eucometis cristata, 454.
cristata affinis, 451, 453.
spodocephala, 451-453.
spodocephala pallida, 451-453.
spodocephala stictothorax,
451-453.

Euetheia canora, 322, 399. Euryzona sepiaria, 196. Evermann. B. W., birds of Carroll County. Indiana, 344. FALCO æsalon, 395. Falco columbarius, 146, 271, 350. columbarius suckleyi, 146. mexicanus, 139. peregrinus anatum, 146. rusticolus, 197. sparverius, 20, 146, 271, 350.

tinnunculus, 110, 205, 395.
Faxon, W., and Allen, J. A., notes on the summer birds of Holderness, Bethlehem, and Franconia, N. H., 149.

Finch, Grass, 18. Pine, 209, 325.

Purple, 12, 150, 152, 209, 273. Rufous-crowned Summer, 40.

Fisher, A. K., Rallus longirostris crepitans breeding on the coast of Louisana, 108; Megascops asio floridanus in Louisana. 111; Quiscalus quiscula aglæus in Louisana, 113.

Fisher, G. J., polydactylism in birds, 218.

Fishhawk, 350.

Flicker. 21, 271. Red-shafted, 16.

Red-shafted, 16. Flycatcher, Acadian, 21, 272.

Crested, 21, 149, 151, 272. Great-crested, 200. Hammond's, 258. Least, 150, 151. Olive-sided, 151, 390. St. Lucas, 90, 396. Slender-billed, 137. Traill's, 151. Wright's, 258.

Yellow-bellied, 151. Fowl, Sage, 367. Fratercula arctica, 134, 135, 171. Fregata aquila, 69, 170, 173, 378. Fringilla canariensis, 413.

gayi, 194. maderensis, 412, 413. morelleti, 413.

Fulica americana, 57, 144, 267, 270,

273, 348. atra, 57. caribæa, 58. martinica, 57. noveboracensis, 56. spinosa, 289, 290, 292, 297,

Fuligula affinis, 65.
americana, 66.
collaris, 66.
ferina var. americana, 66.
marila, 65.
mariloides, 66.

Fuligula rubida, 68. rufitorques, 66. Fulix affinis, 65. collaris, 66.

GADWALL, 173, 268.
Galbraith, C. S., Bachman's Warbler (Helminthophila bachmani) in Louisiana, 323.

in Louisiana, 323.
Galeoscoptes carolinensis, 24, 151, 153, 154, 277, 329, 388, 389, 434.

Gallinago delicata, 144, 270, 328,

330, 331, 348. Gallinula chloropus, 56. galeata, 56, 270, 328, 348. galeata sandvicensis, 197. martinica, 57. Gallinule, Florida, 270, 328, 348.

Gallinule, Florida, 270, 328, 348 Purple, 109, 270. Gannet, 130-135, 278, 282.

Gannet, 130-135, 278, 2 Gavia alba, 202.

Gelochelidon anglica, 76. aranea, 76. cayennensis, 77. nilotica, 76, 374.

nilotica, 76, 374.
Geothlypis agilis, 187, 323, 326.
coryi, 157.
formosa, 24, 422.
macgillivrayi, 35, 362.
philadelphia, 153, 323, 434.
tanneri, 157.
trichas, 24, 151, 153, 154, 276,

326, 389, 434. trichas occidentalis, 35, 362. Glaucidium gnoma, 146, 366, 370. gnoma californicum, 366, 370. gnoma hoskinsii, 136, 372,

> 395. phalænoides, 366, 371.

Glaucion clangula, 422. Glaucionetta clangula americana, 15, 67, 142, 203, 347.

islandica, 16, 142, 347. Gnatcatcher, Blue-gray, 25, 166,

Plumbeous, 166. Gnathospiza raimondi, 412. Godwit, Marbled, 349.

Golden-eye, American, 15, 347, 379. Barrow's, 16, 347.

Goldfinch, 150, 207. American, 22, 152, 273. European, 211.

Goose: Blue 394. Canada, 17, 347. Greater Snow, 183, 379. White-fronted, 143. Gormley, M. H., nesting of the Black Swift, 424.

Goss, N. S., feeding habits of Pelecanus erythrorhynchos, 25; notes on Gymnostinops montezumæ, 27; new and rare birds found breeding on the San Pedro Martir Isle, 240; how far West has Anas obscura been found?, 444.

Gosse, P. H., death of, 446. Grackle, Boat-tailed, 273. Bronzed, 19, 113, 208, 324. Florida, 113, 208, 273.

Purple, 19, 22, 324. Graculus dilophus var. floridanus, 70.

floridanus, 70. mexicanus, 71.

Grassquit. Melodious, 322.

Grebe, 203. Horned, 373. Pied-billed, 268, 345, 373. Western, 15, 201.

Grosbeak, Blue, 22, 119.
Evening, 113, 114, 208, 425.
Kodiak Pine, 397.
Pine. 321.
Rose-breasted, 150, 152.
Western Blue, 399.

Grouse, Canada, 317. Ruffed, 20, 149, 151, 314, 338-340, 349.

Sooty, 367. Grus canadensis, 50, 144. mexicana, 50, 144, 205, 348. poliophæa, 50.

Guara alba, 269. Guillemot, Kittlitz's, 410. Guiraca argentina, 412.

cærulea, 22. cærulea eurhyncha, 119, 399. 412. cyanea, 412.

Gull, American Herring, 345, 374.
Bonaparte's, 345, 374.
Great Black-backed, 107, 318.
Herring, 107.
Ivory, 202.
Laughing, 107, 374, 377.
Little, 170, 171, 393.
Point Barrow, 393.
Ring-billed, 107, 345, 374.

Western, 241. Gymnostinops montezumæ, 27, 99. HABIA ludoviciana, 150, 152, 154, 327, 389. melanocephala, 359, 438-444.

Hæmatopus frazari, 84, 395.

Hæmatopus galapagensis, 85. palliatus, 85, 86.

Haliæetus leucocephalus, 18, 145, 149, 271, 350.

Haliplana anæstheta, 77. discolor, 77.

fuliginosa, 78.

Hancock, J. L., the northern range of *Oporornis formosa* in Illinois, 210; impeded migration and destruction of birds at Chicago, 432.

Hardy, F. P., testimony of some early voyagers on the Great Auk, 380.

Harporhynchus, bendirei, 116, 117, 118, 161. crissalis, 161.

curvirostris palmeri, 116, 161, rufus, 16, 24, 151, 154, 277, 325, 388, 389, 434.

Harrier, Marsh, 350.
Harris, G. E., early nesting of Otocoris alpestris praticola, 320.
Hasbrouck, E. M., Dendroica cærulea in the District of Colum-

bia, 323. Hawk, American Sparrow, 20, 271. Broad-winged, 20, 350. Cooper's, 20, 246, 271, 316, 350. Hen, 149. Little Black, 185.

Little Black, 185.
Marsh, 151, 271.
Pigeon, 271, 350.
Red-shouldered, 20, 271, 350.
Red-tailed, 18, 271, 350.
Rough-legged, 350.
Sharp-shinned, 20, 271, 350.
Sparrow, 350.
Swainson's, 424.

Western Red-tailed, 205. Helinaia swainsonii, 187, 323, 325. Helminthophila bachmanii, 187,

110thophila bachmanii, 187, 323, 325, 428. celata, 275, 323. celata lutescens, 34, 360, 361. chrysoptera, 391. cincinnatiensis, 401. lawrencei, 401, 421, 427. leucobronchialis, 115, 148, 401, 421, 427. luciæ, 33. peregrina, 152, 187, 326. pinus, 24, 427. ruficapilla, 150, 152, 323, 326, 389. ruficapilla gutturalis, 33, 360.

virginiæ, 33.

Helmitherus vermivorus, 23, 275. Heron. Bahama Green, 83. Black-crowned Night, 20

Black-crowned 268, 269, 348. Frazar's Green, 83, 394. Great Blue, 268, 269, 347. Green, 19, 269, 348. Little Blue, 268, 269. Louisiana, 268, 269. Night, 327, 328. Snowy, 269. Ward's, 183.

Wuerdemann's, 195, 394. Hesperocichla nævia, 365. Heteractitis incanus, 237. Himantopus knudseni, 197. Himatione parva, 197. Holder, J. B., death of. 220.

Hopkins, C. L., notes relative to the sense of smell in the Turkey Buzzard (*Carthartes aura*), 248. Hudson, W. H., see Sclater, P. L. Hummingbird, Floresi's, 396.

Ruby-throated, 21, 151, 272. Hydralector cristatus, 299, 302. gallinaceus, 290, 292, 293, 302.

novæ-hollandiæ, 303. Hydrochelidon fissipes, 79. fuliginosa, 78. lariformis, 79.

nigra, 79. nigra surinamensis, 79, 140, 376. plumbea, 79.

surinam, 79. Hydrophasianus chirurgus, 289, 291, 292, 293, 294, 303, 304. sinensis, 304.

IBIS, White, 269. Icteria virens, 24, 116, 389, 392, 405. virens longicauda, 36, 438,

virens longicauda, 36, 43 441-444. Icterus bairdi, 158.

bullocki, 119, 262. galbula, 22, 150, 152, 154, 273, 388, 389. gularis, 454-456. gularis, vucatanensis, 454-456.

gularis yucatanensis, 454-456. spurius, 22, 119, 186, 211, 273, 413. wagleri, 92.

wagleri castaneopectus, 91. Ictinia mississippiensis, 270. Indigo-bird, 150.

Ionornis martinica. 57, 109, 270. JACANA gymnostoma, 402. indica, 292.

jacana, 291, 292, 294, 295.

Jacana nigra, 292, 294, 295, 296, 297, spinosa, 289, 290, 292, 294, 297, 402, violacea, 51.

Jay, Black-headed, 396.
Blue. 21, 150, 151, 434.
Canada, 316.
Florida Blue, 112, 272, 324,
Oregon, 226.
Xantus's, 397.

Jeffries, J. A., a description of an apparently new species of *Tro-chilus* from California, 168.

Johnson, H. P., the Yellow-breasted Chat breeding in Malden, Mass., 116.

Joree, 274.
Junco hyemalis, 149, 150, 152, 154, 325, 388, 389, 391.
hyemalis carolinensis, 398, hyemalis oregonus, 359.
hyemalis shufeldti, 398.

Junco, 325, 388. Carolina, 398. KESTREL, 395.

European, 110, 205.
Keyes, C. R., occurrence of Coccothraustus vespertina in Iowa, 114: Blackbird flights at Burlington, Iowa, 207; Dendroica tigrina at Iowa City in November,

na at Iowa City in November 211.
Killdeer, 20, 270, 349.
Kingbird, 21, 149, 151, 200, 272.
Kingfisher, 149.

Belted, 21, 151, 271. Kinglet, Golden-crested, 15, 337, 388. Golden-crowned, 153.

Ruby-crowned, 153. Ruby-crowned, 15, 166, 277. Kite, Mississippi, 270.

Swallow-tailed, 270. Kittiwake, 134.

Knot, 110. Kumlien, F., reason or instinct?,

Kumlien, L., observations on bird

migration at Milwaukee, 325; see also Kumlien, F., 434-435. LAGOPUS rupestris atkhensis, 410.

Lanius ludovicianus, 275, 312. ludovicianus excubitorides, 31, 360. ludovicianus gambeli, 360,

299. Lark, Horned, 181. Prairie Horned, 111, 170, 180, 206.

Larus affinis, 197.

Larus argentatus, 75.
argentatus smithsonianus,
107, 345, 374.
argentatus var. vegæ, 310.
atricilla, 75, 107, 374, 377.
barrovianus, 393.
cachinnans, 310, 402.
delawarensis, 107, 345, 374.
franklinii, 75.
marinus, 75, 107, 318.
minutus, 170, 171, 393.
occidentalis, 241.

philadelphia, 76, 171, 345, 374; schistisagus, 197, smithsonianus, 75, vegæ, 310. zonorhynchus, 75.

Lateriallus gossei, 54.

Lawrence, G. N., description of a new species of Wren from the Island of Tobago, West Indies,

Leucosticte griseinucha, 410. Leucosticte, Aleutian, 410. Limnopardalus variegatus, 52. Limosa fedoa, 349. Limosa melanuroides, 197.

Linden, C., death of, 220. Loggerhead, 275. Longspur, Chestnut-collared, 14,

Lapland, 181. McCown's 14, 321.

Loomis, L. M., notice of the presence of Quiscalus quiscula æneus in upper South Carolina, 113; on the further occurrence of Otocoris alpestris praticola in Chester County, South Carolina, 206; Seiurus noveboracensis notabilis in South Carolina, 324.

Loon, 149, 268, 345, 373, 390. Red-throated, 147, 149, 170, 374. Sheldrake, 171. Lophodytes cucullatus, 69, 141, 268,

346, 378. Loxia curvirostra minor, 22, 115, 152, 154, 208, 317, 324, 325, 340, 389, 390.

curvirostra stricklandi, 114, 358. leucoptera, 152, 410.

Loxigilla barbadensis, 158.
richardsoni, 158.
Lucas, F. A., the Bird Rocks of
Gulf of St. Lawrence in 1

Lucas, F. A., the Bird Rocks of the Gulf of St. Lawrence in 1887, 129; notice of his 'Notes on the Osteology of the Spotted Tinamou (Nothura maculosa), 198; Great Auk notes, 278; abnormalities in the ribs of birds, 329. Lunda cirrhata, 328. MACRORHAMPHUS griseus, 110.

scolopaceus, 147, 184.
Magpie, American, 317.
Mallard, 17, 268, 313, 314, 346, 378.
Mareca americana, 63.
Margarops montanus albiventris,

montanus rufus, 47, 157. Martin. Cuban, 399. Purple, 23, 31, 150, 152, 275. McIlwraith, K. C., Prothonotary

McIlwraith, K. C., Prothonotar Warbler in Ontario, 322. Meadowlark, 15, 22, 388. Mexican, 273.

Western, 413.
Megalopterus stolidus, 80.
Megascops asio, 20, 351, 389.
asio floridanus, 111, 271, 324.
asio kennicottii, 146, 369.
asperus, 87.
hastatus, 194.
vermiculatus, 194.

vinaceus, 88.
Melanerpes carolinus, 21, 271.
erythrocephalus, 21, 271, 433.
formicivorus bairdi, 240.
torquatus, 240, 255.

Melanoptila glabrirostris, 329. Melagris gallopavo, 18, 204, 27c, 314, 349.

Melospiza fasciata, 22, 150, 152, 154, 211, 274, 326, 389. fasciata guttata, 359. fasciata heermanni, 359. fasciata montana, 17, 139, 359. georgiana, 150, 152, 274, 327, 328, 389. lincolni, 18, 148, 359.

Merganser americanus, 141, 346. serrator, 141, 346, 378.

Merganser, American, 346. Hooded, 268, 346, 378. Red-breasted, 346, 378.

Mergus albellus, 422. anotarius, 422. cucullatus, 69.

Merlin, 395.
Merriam, C. H., what birds indicate proximity to water, and at what distance, 119; Euetheia canora from Sombrero Key, Florida, 322; description of the breeding

15

0

plumage of Chadbourne's Field Sparrow (*Spizella arenacea*), with evidence of its specific distinctness, 402.

tinctness, 402.

Merrill, J. C., notes on the birds of Fort Klamath, Oregon, with remarks on certain species by William Brewster, 139, 251, 357.

Merula migratoria, 25, 151, 153, 155, 211, 277, 316, 327, 389, 434

migratoria propinqua, 167, 365.

Metopidius æneus, 299. africanus, 289, 292, 293, 301. albinucha, 293, 300. gallinaceus, 293.

gallinaceus, 293. indicus, 293, 299. œnea, 289. Micrathene whitneyi, 372.

Micropalama himantopus, 110.
Miller, G. S. Jr., description of an apparently new *Poocates* from Oregon, 404.

Milvulus tyrannus, 158. Mimocichla ravida, 156. Mimus caudatus, 160.

polyglottos, 24, 119, 159, 183, 276, 314, 421.

Mitrephanes phæocercus, 137. phæocercus tenuirostris, 137. Mniotilta varia, 23, 150, 152, 275,

Mniotilta varia, 23, 150, 152, 275, 326, 389, 433, 434. Mockingbird, 24, 119, 159, 183, 276,

Molothrus ater, 15, 21, 151, 207,

Moore, H. F., Protonotaria citrea in Montgomery County, Penn-

sylvania, 210. Morris, R. O., Larus atricilla at Springfield, Mass., 107.

Motacilla ocularis, 410. Murre, 131-135, 281, 382. Muscisaxicola occipitalis, 194. Myadestes townsendii, 167, 365.

Myiarchus berlepschii, 266. crinitus, 21, 149, 151, 154, 272, 389.

cristatus, 200. denigratus, 158. Myiobius ridgwayi, 457. xanthopygus, 457.

Myiodynastes luteiventris, 385. Nannocnus, 196.

Nettion carolinensis, 64. Nighthawk, 21, 149, 151, 314. Chapman's, 186, 272. Nomonyx dominicus, 68. Nonpareil, 119.

Norton, R., Ceophlaus pileatus in Franklin County, Massachusetts, 111.

Notherodius scolopaceus, 51. Northura maculosa, 106, 198. Nucifraga caryocatactes leptorhyn-

chus, 422. caryocatactes pachyrhynchus, 422.

Numenius cyanopus, 197. longirostris, 14, 145.

Nuthatch, Brown-headed, 277, 432.
Pygmy, 165, 362.
Red-bellied, 118, 393.
Red-breasted, 153, 165.
Slender-billed, 165.

White-breasted, 25, 153, 277. Nyctala acadica, 146, 351, 369, 389, 390.

Nyctea nyctea, 315, 351. Nyctherodius violaceus, 48.

Nyctiardea gardeni, 49. grisea nævia, 49. violacea, 48.

Nycticorax americanus, 49. gardeni, 49.

nycticorax nævius, 20, 49, 144, 198, 269, 327, 328, 348. violaceus, 48. vulgaris, 49.

Nyctidromus, 205.

Nyctidromus albicollis derbyanus, 46.

albicollis merrilli, 44, 402. Nyroca affinis, 66.

americana, 66. collaris, 66. ferruginea, 64. vallisneria, 66.

OCEANITES oceanicus, 80. wilsoni, 80.

Oceanodroma hornbyi, 401. leucorhoa, 132, 280. leucorhoa macrodactyla, 394.

Estrelata caribæa, 81. hesitata, 81. jamaicensis, 81.

Old square 202,

Old-squaw, 203. Old-wife, 203.

Olor buccinator, 143, 347. columbianus, 143, 176, 347. Onychoprion fuliginosus, 78.

Oporornis formosa, 210.

Oreomyza bairdi, 197. Oriole, Baltimore, 22, 150, 152, 273, 388. Bullock's, 119.

Chestnut-breasted, 91. Orchard, 22, 119, 186, 273.

Oropendula, 27. Oroscoptes montanus, 159. Ortygometra carolina, 55. flaviventris, 54. jamaicensis, 55.

minuta, 54. Osprey, 350.

Osprey, American, 271. Otocoris alpestris, 111, 148, 181, 442, 443.

alpestris arenicola, 15, 259, 260. alpestris practicola, 111, 148,

170, 180, 206, 320. alpestris strigata, 259. Otophanes mcleodii, 89.

Oven-bird, 24, 150, 153, 276, 338, 339.

Owl, Acadian, 369. American Barn, 20, 180, 271. American Long-eared, 20. Barred, 20, 248, 350. California Pygmy, 366. Chihuahua Screech, 88. Elf. 372.

Ferruginous Pygmy, 366, 371. Florida Barred, 271, 324. Florida Screech, 111, 271, 324.

Great Gray, 110. Great Horned, 18, 271, 313, 351.

Hawk, 410. Hoskins' Pygmy, 136, 372,

Kennicott's, 369. Lapp, 410. Long-eared, 350. Pygmv, 366. Saw-whet, 351. Screech, 20, 351. Short-eared, 350. Snowy, 315, 351. Spotted Screech, 87.

Oyster-catcher, Frazar's, 84, 395.

PALMEN, J. A., notice of his con-tributions to the knowledge of the bird fauna of the Siberian coasts of the Arctic Sea, 306.

Palmer, W., see Smith, H. M. Pandion haliaëtus carolinensis, 146, 271, 350.

Parauque, Merrill's, 44. Pardirallus variegatus, 52. Park, A. F., breeding of the Prairie Horned Lark in Eastern New York-a correction, 111. Paroquet, Carolina, 148, 185. Parra ænea, 299.

> albinucha, 289, 290, 300. atrata, 299. atricollis, 300. brasiliensis, 289, 290, 296. capensis, 290, 292, 301. chilensis, 290. chirurgus, 289. cordifera, 289, 290, 297. cristata, 289, 290, 299, 303. cuprea, 299. gallinaceus, 289, 290, 292, 302. gymnostoma, 51, 289, 290, 297, 298. hypomelæna, 290, 296.

africana, 289, 290, 292, 301.

indica, 289, 290, 299. intermedia, 290, 294. jacana, 51, 289, 290, 292, 294, 295, 296, 297, 298. luzoniensis, 289, 304. melanochloris, 299.

melanopygia, 290, 296, 297. melanoviridis, 299. nigra, 289, 290, 296, 297. novæguineæ, 290, 303. œnea, 289. phænicura, 304. sinensis, 289, 304.

spinosa, 289, 290. superciliosa, 299. variabilis, 289, 297, 298, 299. violacea, 51, 290, 297, 299. viridis, 289, 290, 296. Parrot, Thick-billed, 395.

Parus atricapillus, 151, 153, 155, 363, 389.

atricapillus septentrionalis, 119. atricristatus, 119. castaneifrons,

atricristatus bicolor, 25, 277, 421. bicolor texensis, 400 carolinensis, 25, 277 carolinensis agilis, 46, 400.

cinctus obtectus, 410. gambeli, 118, 359, 362, 363. hudsonicus, 153, 410. rufescens, 363. stoneyi, 400.

wollweberi, 166.

Passer domesticus, 22, 152, 200, 312, 314, 315, 412. griseigularis, 412. montanus saturatus, 196. Passerella iliaca, 18. iliaca megarhyncha, 359. iliaca unalaschcensis, 359. Passerina amœna, 359. ciris, 119. cyanea, 22, 150, 152, 154, 274, 389. versicolor pulchra, 399. Pediocætes phasianellus columbianus, 145. Pelecanus aquilus, 69. erythrorhynchos, 25, 99, 141,

147, 346, 377.
fuscus, 25, 70, 377.
leucogastra, 72.
piscator, 72.
sula, 72.
Pelican, American White, 346, 377.
Brown, 25, 70, 377.

Frigate, 173.
White, 25, 96.
Peocilonetta bahamensis, 62.
Perisoreus canadensis, 316.
obscurus, 226, 261.

Petrel, Guadaloupe, 394. Leach's 132, 280. Petrochelidon lunifrons. 23, 31 150, 152, 154, 360, 389.

Peucæa æstivalis, 274, 355.
æstivalis bachmanii, 274, 351.
arizonæ, 401.
homochlamys, 398, 412.
mexicanus, 401.
ruficeps, 40, 41.
ruficeps boucardi, 40, 41, 42,
401.
ruficeps eremæca, 40, 41, 401.

ruficeps scottii, 41,42, 398, 412.

ruficeps var. boucardi, 412. Pewee, Richardson's, 258. Wood, 21, 150, 151, 272. Phacellodomus inornatus, 194. Phæornis myiadestina, 197. Phæospiza thomensis, 317.

Phæospiza thomensis, 317. Phaëthon æthereus, 74. flavirostris, 73.

Phaëton æthereus, 74, 241, 244. flavirostris, 73. Phainopepla nitens, 31.

Phainopepla, 31
Phalacrocorax dilophus cincinatus,
141.
dilophus floridanus, 70, 377.
floridanus, 70.

Phalacrocorax mexicanus, 71.
perspicillatus, 401.
resplendens, 71.
townsendi, 71.
Phalænoptilus nuttalli californicus.

396. nuttalli nitidus, 396. Phalarope, Wilson's, 177. Phalaropus tricolor, 144, 177. Phasianurus vigorsii, 62. Pheasant, Golden, 315. Philacte canagica, 311.

Philohela minor, 20, 270, 330, 331, 348, 389. Phæbe, 21, 150, 151, 272.

Phoys, 196.
Phrygilus aldunatii, 412.
atriceps, 194, 412.
formosus, 194.
gayi, 194.
punensis, 194, 412.

saturatus, 412. Pica pica hudsonica, 261, 317, 414. Picicorvus columbianus, 262. Picoides arcticus, 151, 240, 254, 390. Picolaptes rikeri, 194.

Picumnus minutus, 212.
Picus auratus, 212.
bengalensis, 212.
carolinus, 212.
luridus, 212, 214.
macci, 212.
martius, 212.
medius, 212.
tridactylus, 212.

villosus harrisii, 228, 254. viridis, 214. Pigeon. Passenger, 270, 349. Viosca's, 86, 395.

White-crowned, 185.
White-growned, 185.
Wild, 314, 349.
Pindar, L. O., Pine Grosbeak in

Fulton Co., Kentucky, 321. Pinicola enucleator, 321. enucleator kodiaka, 397.

Pintail, 269, 346, 378. Pipilo chlorurus, 359, 412, 438-444erythrophthalmus, 22, 150, 207, 274, 388, 389. erythrophthalmus alleni, 274. maculatus arcticus, 16. maculatus megalonyx, 139, 359, 438-444-

maculatus oregonus, 359. Pipit, American, 159, 276. Meadow, 410.

Piranga erythromelas, 23, 30, 150, 152, 181, 322, 327, 389, 433

Piranga hepatica, 30.
ludoviciana, 29, 359, 438, 441444rubra, 23, 30, 181, 274rubra cooperi, 30.
rubriceps, 399.
Platypsaris albiventris, 425.
Platyrhynchus xanthopygus, 457.
Plautus impennis, 278, 331, 380, 422.
Plectrophenax nivalis townsendi, 397.
Plotus anhinga, 71.

Plotus anhinga, 71.
Plover. Black-bellied, 110.
Golden, 308.
Snowy, 110.
Podiceps dominicus, 155.
holbolli, 156.

Podilymbus carolinensis, 156. dominicus, 155. podiceps, 140, 156, 268, 345,

Polioptila cærulea. 25, 166, 277, 3²⁴. cærulea cæsiogaster, 157. plumbea, 166.

Poocetes gramineus, 22, 150, 152, 154, 273, 389. gramineus affinis, 404. gramineus confinis, 18, 358.

Poor-will, Frosted, 396. Poospiza boliviana, 412. Porphyrio martinica, 57. Porphyrospiza pulchra, 412. Porphyrula martinica, 57.

Porzana alfari, 194.
carolina, 20, 38, 39, 55, 144,
269, 328, 348, 414, 422.
concolor, 54.
flaviventer, 55.
flaviventris, 54, 55.
jamaicensis, 55.
minuta, 55.
noveboracensis, 56, 147, 177,

phæopyga, 196.
Price, W. W., occurrence of Vireo flavoviridis at Riverside, California, 210; nesting of the Red-faced Warbler (Cardellina rubrifrons) in the Huachuca Mountains, Southern Arizona, 385; Xantus's Becard (Platypsaris albiventris) in the Huachuca Mountains, Southern Arizona, 425.

Procellaria diabolica, 81. hasitata, 81. jamaicensis, 81. obscura, 82.

319

Procellaria oceanica, 80.
Progne cryptoleuca, 399.
subis, 23, 31, 150, 152, 275, 389.
Protonotaria citrea, 170, 182, 210,

275, 322, 325. Pryer, H. J. S., death of, 332. Psaltriparus lloydi, 43, 400. melanotis, 44. minimus, 364.

minimus, 364. minimus californicus, 364. plumbeus, 44, 166, 364. Pseudochloris mendozæ, 412.

Psittacula cyanochlora. 461, 462. deliciosa, 461, 462. exquisita, 461, 462. guianensis, 462. passerina, 460, 462. passerina vivida, 460. spengeli, 461, 462.

Psittirostra psittacea, 197. Ptarmigan, Turner's, 410. Pterocyanea discors, 63. Pterodroma caribbæa, 81. Publications received, 106, 200, 317, 422.

Puffin, 134, 135, 171, 328. Puffinus auduboni, St. 82, 173. borealis, 108, 170, 173, 202. major, St. obscurus, St.

major, 81. obscurus, 81. stricklandi, 81. Pyrrhospiza humii, 412. punicea, 412.

Pyrrhula cassini, 197, 410. cineracea. 197. griseiventris, 197. kurilensis, 197. pyrrhula europæa, 197. pyrrhula kamtschatica, 197. rosacea, 197.

Pyrrhuloxia sinuata, 119. sinuata beckhami, 398. sinuata peninsulæ, 398. Pyrrhuloxia, Arizona, 398.

St. Lucas, 398.
QUAIL. Messina, 314.
Querquedula carolinensis, 64.
clypeata, 65.
cvanoptera, 64.
discors, 63.

inornata, 64.

Quiscalus caymanensis, 158.
major. 273.
purpureus, 329.
quiscula. 19. 22. 113. 208. 324.
quiscula æneus. 15, 19, 113,
207. 208, 324. 389.
quiscula aglæus, 113, 208, 273.

RAIL, Carolina, 38, 39, 328, 414. King, 20, 176, 269. Scott's, 305, 394. Virginia, 20. Yellow, 147, 177, 319.

Rallina castanea, 54. Rallus carolinus, 55.

concolor, 54. corvi, 53. crepitans, 53.

elegans, 20, 52, 53, 176, 269. flaviventer, 54.

giganteus, 50. jamaicensis, 55. lariformis, 79. longirostris, 53

longirostris caribæus, 53, 319, 394.

longirostris crepitans, 53. longirostris scottii, 305, 394. longirostris var. caribæus, 53.

maculatus, 52. minutus, 54. superciliaris, 54. variegatus, 52. virginianus, 53.

Raven, 329. Mexican, 241. Northern, 396.

Razorbill, 134, 135, 382. Recurvirostra americana, 16.

Redstart, 151. American, 24, 36, 153, 276, 433. Painted, 36.

Redwing, Bahaman, 397. Sonoran, 397

Reed, J. H., an albino Blackpoll Warbler, 432.

Regulus calendula, 15, 166, 277, 365.

cuvieri, 401. satrapa, 15, 153, 326, 337, 388, 389, 393, 433, satrapa olivaceus, 364.

Rhamphastos cuvieri. 122. Rhamphocinclus sanctæ-luciæ, 157. Rhoads, S. N. occurrence of Larus marinus on the southern coast of

New Jersey, 318. Rhodospingus cruentus, 411. mentalis, 412.

Rhodospiza obsoleta, 411. Rhyncaspis clypeata, 64. Rhynchophanes mccownii, 14, 321.

Rhynchopsitta pachyrhyncha, 395. Richmond, C. W., an annotated list

of birds breeding in the District of Columbia, 18.

Ridgway, R., Spencer Fullerton Baird, 1; notice of his descriptions of new American birds, 194; notice of his 'Notes on Ardea wuerdemanni', 195; notice of his paper on 'Trogon ambiguus breeding in Arizona,' 195; description of a new Tityra from western Mexico, 263; supplementary remarks on the genus Psittacula Brisson, 460.

Rissa pollicaris, 310. tridactyla, 134, 310. Rives, W. C., Jr., Cory's Shearwater at Newport, R. I., 108.

Robin, 151, 316, 327. American, 25, 153, 277. Western, 167.

Roddy, H. J., feeding habits of some young Raptores, 244. Rostratula, 204. Rufirallus concolor, 54.

Rynchops nigra, 74, 376. SALPINCTES obsoletus. 162.

Sanderling. 179. Sandpiper, Bartramian, 270.

Least, 349. Pectoral, 349. Purple, 178, 184. Semipalmated, 349. Solitary, 270, 330, 349. Spotted, 20, 149, 151, 270, 349. Stilt, 110.

Western, 184. Sapsucker, Red-breasted, 229. Red-naped, 226. Williamson's 235

Yellow-bellied, 225, 271, 430 Saunders, W. E., first occurrence of the Western Red-tail in Ontario, 205; the Evening Grosbeak in Ontario, 208.

Sayornis phæbe, 21. 150. 151. 272. Schistospiza griseocristata, 411. Sclater, P. L., and Hudson, W. H., notice of their 'Argentine Ornithology,' 199.

Scolecophagus carolinus, 207, 273. cvanocephalus, 15, 262.

Scoter, Surf. 202, 203. Scott, W. E. D., on the avifauna of Pinal County with remarks on some birds of Pima and Gila Counties, Arizona, with annotations by J. A. Allen, 29, 159;

supplementary notes from the Gulf Coast of Florida, with a description of a new species of Marsh Wren, 183; a summary of observations on the birds of the Gulf Coast of Florida, 373; Bachman's Warbler (Helminthophila bachmani) at Key West, Florida, in July and August, 428; Dendroica coronata at Key West in summer, 430.

summer, 430.
Seebohm. H., notice of his 'Geographical Distribution of the Charadriidæ', 189.

Seiurus aurocapillus, 24, 150, 153, 154, 276, 315, 326, 327, 338, 380, 434.

389, 434. motacilla, 24, 276, 421. noveboracensis, 150, 153, 326, 327, 434.

noveboracensis notabilis, 35, 148, 324.

Sennett, G. B., notes on the *Peucœu ruficeps* group, with description of a new subspecies, 40; descriptions of a new species and two new subspecies of birds from Texas, 43; unexpected occurrence of certain shore birds in Texas in midsummer and in breeding plumage, 110; dichromatism in the genus *Nyctidromus*, 205; a new form of Clapper Rail, 305; an addition to the list of North American birds, 319. Setophaga picta, 36.

ruticilla, 24, 36, 151, 153, 154,

276, 326, 389, 433.
Sharpe, R. B., notice of his catalogue of the family Fringillidæ,

Shearwater, Audubon's, 173. Cory's, 108, 170, 173, 202.

Shorelark, 15. Shoveller, 269, 346.

Shrike. Loggerhead, 275, 312. White-rumped, 31.

White-rumped, 31.
Shufeldt, R. W., notice of his 'Review of the muscles used in the classification of birds,' 104; notice of his report on birds' sterna and skulls collected by Dr. Streets, 198; observations on the pterylosis of certain Picidæ, 212; notes on Melanerpes torquatus, 319; osteological notes upon Puffins and Ravens, 328; the sternum in the Solitary Sandpiper and

other notes, 330; notice of his paper on the osteology of the Icteridæ and Corvidæ, 413; notice of his 'Osteology of Porzana carolina,' 414; the osteology of Habia melanocephala with comparative notes upon the skeletons of certain other conirostral birds and of Tanagers, 438.

Sialia arctica, 168, 366. mexicana, 167, 365, 366. sialis, 22, 25, 151, 153, 155, 277, 389, 433.

Simorhynchus pygmæus, 410. Siskin, Pine, 152, 426.

Sitta canadensis, 118, 153, 165, 363, 389, 393. carolinensis, 25, 153, 277. carolinensis aculeata, 165,

363. pusilla, 277, 432. pygmæa, 165, 362, 363.

Skimmer, Black, 376. Skylark, 180.

Missouri, 15.
Slade, D. D., the Wild Turkey in Massachusetts, 204.

Smith, H. M., and Palmer, W., additions to the avifauna of Washington and vicinity, 147.

Smythe, E. A., Jr., the Old-squaw (Clangula hyemalis) in South Carolina, 203.

Snipe, 331. English, 328.

Wilson's, 270, 330, 348. Snow, F. H., notice of his description of a new ornithichnite, 105; Æchmophorus occidentalis in

Kansas, 201. Snowbird, 149, 150, 152. Snowflake, Prybilof, 397. Solitaire, Townsend's, 167. Somateria spectabilis, 174.

Sora, 20, 269, 348. Sparrow, Acadian Sharp-tailed, 397. Bachman's, 274, 351.

Bay-winged, 152.
Boucard's, 42.
Cahoon's, 93.
Chadbourne's Field, 402.
Chipping, 22, 150, 152, 274.
English, 315.
European, 200.
European House, 22.
Field, 22, 150, 152, 274.

Fox-colored, 18. Grasshopper, 22, 273. Sparrow, Henslow's, 22, 210.

House, 152.

Intermediate White-crowned,

17.

Leconte's, 210.

Lincoln's, 18.

McLeod's, 92.

Pine-woods, 274.

Plateau, 265.

Rufous-crowned, 41.

Savanna, 150, 152, 273.

Scott's, 42, 398. Scott's Seaside, 284, 287, 398.

Song, 17, 22, 150, 152, 274.

Swamp, 150, 152, 274.

Texan Seaside, 286, 398.

Texas, 352. Vesper, 22, 273.

Western Field, 398. Western Tree, 367. Western Yellow-winged, 15.

White-crowned, 181.

White-throated, 17, 149, 150,

152, 274, 388, 426. Spatula clypeata, 17, 64, 65, 142,

269, 346. Spermophila albitorquis, 411.

colombiana, 412.

cucullata, 412.

gutturalis, 158.

morelleti. 412.

morelleti parva, 412.

parva, 402. 412.

plumbea, 412.

polionota, 412.

whitelevana. 412. Sphyrapicus ruber, 226, 229, 236,

240, 255.

thyroideus. 228, 235, 240, 254,

255.

varius, 151, 154, 225, 240, 271,

325, 430.

varius nuchalis, 214, 216, 226,

240, 255, 332. williamsoni. 235.

Spindalis salvini. 157

zena townsendi, 157, 194.

Spinus notatus, 412.

pinus, 152, 209, 325, 358, 426.

psaltria, 358.

tristis, 22, 150, 152, 154, 207,

273. 358, 389.

Spiza americana. 23.

townsendi, 401.

Spizella arenacea, 402.

breweri, 359.

monticola ochracea, 367.

pallida. 311.

Spizella pusilla, 22, 150, 152, 274, 389.

pusilla arenacea, 398.

socialis, 22, 150, 152, 154, 274, 389.

socialis arizonæ, 358.

Spoonbill, Roseate, 269. Sporophila morelleti, 402.

Stejneger, L., propatagialis cucullaris, 120; notice of his papers on Japanese birds, 195; notice of his papers on Hawaiian birds, 196; notice of his 'Catalogue of the Birds inhabiting the Commander Islands.' 197; notice of his 'Notes on the Northern Palæarctic Bullfinches', 197; Palmén's contributions to the knowledge of the bird fauna of the Siberian coasts of the Arctic Sea, 306.

Stelgidopteryx serripennis, 23, 31, 360.

Stercorarius parasiticus, 197.

Sterna acuflavida, 77.

anglica, 76. anosthæta, 77

(Anous) stolida. 80.

antillarum, 79, 141, 375, 376.

argentea, 79.

cantiaca, 77.

cayennensis, 76.

dougalli, 78, 375.

forsteri, 375, 376. fuliginosa, 78.

hirundo, 77, 346, 375, 376. maxima, 76, 374, 376.

minuta, 79.

minuta americana, 79.

nigra, 79. nilotica, 76.

paradisea, 78.

regia, 76.

sandvicensis acuflavida, 77,

374, 376. stolida. 80.

superciliaris, 79.

surinamensis, 79.

wilsoni, 77. Sternula antillarum, 79.

frenata, 79.

Stint, Long-toed, 394. Stone, W.. Helminthophila leuco-

bronchialis in Pennsylvania, 115; Sylvania mitrata at German-

town, Pennsylvania, in November, 116; an abnormal Scarlet

Tanager, 322.

Strix pratincola, 20, 180, 271.

Sturnella magna, 22, 211, 388, 389. magna mexicana, 273. magna neglecta, 15, 262, 329,

413, 414.
Sula bassana, 130, 131, 132, 133, 134, 135, 278, 282.
brewsteri, 241, 242.
cyanops, 71.
dactylatra, 71.
fiber, 72.
fusca, 72.

gossi. 241. leucogastra, 72. parva, 72. piscator, 72, 73. sula, 72. Surnia ulula, 410.

ulula caparoch, 390.
Swallow, Bank, 23, 31, 152.
Barn, 23, 31, 150, 152.
Cliff, 23, 31, 150.
Eave, 152.
Rough-winged, 23, 31.
Tree, 19, 31, 275.
Violet-green, 16, 31.

White-bellied, 16, 150, 152, 433. Swan, Trumpeter, 347. Whistling, 176, 347.

Swift, Black, 424. Chimney, 21, 149, 151, 272. Vaux's, 256.

Swift, E., capture of the Brownheaded Nuthatch near Elmira, N. Y., 432.

Swinburne, J., breeding of the Evening Grosbeak (Coccothraustes vespertina) in the White Mountains of Arizona, 113; occurrence of the Chestnut-collared Longspur (Calcarius ornatus) and also of McCown's Longspur (Rhywcophanes mccownii) in Apache Co., Arizona, 321.

Sycalis brasiliensis, 412. jamaicæ, 412. raimondi, 412. taczanowskii, 412. Sylbeocyclus carolinensis, 156.

dominicus, 155.
Sylvania canadensis, 151. 153, 389, 392, 393.
microcephala, 401.
mitrata, 116, 276.
pusilla, 323, 433.
pusilla pileolata, 36, 362.

Symphemia semipalmata, 185.

Symphemia semipalmata inornata, 145, 185, 395, 423. semipalmata speculifera, 424.

speculifera, 424. Synallaxis coryi, 458. fuliginosa, 459.

Synthliborhamphus antiquus, 401. wumizusume, 401.

Syrnium maingayi, 201. nebulosum, 20, 248, 350, 389. nebulosum alleni, 271, 324.

TACHYCINETA bicolor, 16, 19, 31, 150, 152, 275, 360, 389, 433, 434.

thalassina, 16, 31, 360. Tachypetes aquilus, 69, 319. Tanager, Cooper's, 30.

Hepatic, 30. Louisiana, 29. Scarlet, 23, 150, 152, 181, 322, 433. Summer, 23, 181, 274.

Tattler, Wandering, 237.
Teal, Blue-winged, 17, 269, 346, 378.
Green-winged, 17, 269, 3 4,

Tern, Black, 376.
Cabot's, 374, 376.
Common, 346, 375, 376.
Forster's, 375, 376.
Gull-billed, 374.
Least, 375.
Roseate, 375.

Royal, 374. Thalasseus acuflavidus, 77. cayanus, 76. regius, 76.

Thalassidroma wilsonii, 80. Thalassogeron culminatus, 107, 318. Thaumatias cæruleiceps, 264. Thorne, P. M., abnormal, plumage

Thorne, P. M., abnormal plumage of Xanthocephalus xanthocephalus, 112.

Thrasher, 277.

Bendire's, 116, 117, 118, 161.

Brown, 24, 151, 277, 388.

Crissal, 161.

Palmer's 116, 161.

Sage, 159.

Thrush, Audubon's Hermit, 167.

Bicknell's, 153.

Brown, 16, 325.

Dwarf Hermit, 167.

Gray-cheeked, 277.

Hermit, 151, 153, 277, 338.

Olive-backed, 153.

Thrush, Willow, 167. Wood, 25.

Thryothorus bewickii, 324. bewickii bairdi, 402. bewickii murinus, 163, 402. guadeloupensis, 157.

ludovicianus, 24, 183, 277, 421. ludovicianus berlandieri, 277.

ludovicianus miamensis, 187. Thurber, E. C., notice of his 'List of Birds of Morris County, New Jersey,' 421.

Tichodroma muraria, 423. Tinamou, Spotted, 106, 198. Titmouse, Black-capped, 119. Bridled, 166. Chestnut-fronted, 400. Hudsonian, 153. Mountain, 359, 362. Texan Tufted, 400.

Tufted, 25, 277 Tityra albitorques, 287. nigriceps, 287. personata, 287

personata griseiceps, 263. Torrey, B., a second instance of the breeding of the White-throated Sparrow in Eastern Massachusetts, 426.

Totanus crassirostris, 424. flavipes, 270, 327, 330, 331, 332, 349. melanoleucus, 110, 145. 327, 330, 331, 349. ochropus, 330, 332. solitarius, 144, 270, 330, 349.

speculiferus, 423. Towhee, 16, 22, 150, 207, 274, 388. White-eyed, 274.

Townsend, C. H., notice of his 'Field-notes on the mammals, birds, and reptiles of Northern

California', 104.
Treat, W. E., Helminthophila celata in Connecticut, 323.

Tringa alpina, 307. alpina pacifica, 307. canutus, 110. chirurgus, 289, 292, 303. cooperi, 401. damacensis, 394. maculata, 349. maritima, 178, 184. minutilla, 308, 349, 366. ruficollis, 308.

temminckii, 308. Trochilus alexandri, 169. anna, 168.

Trochilus calliope, 257. colubris, 21, 151, 154, 272, 329. floresii, 396. lucifer, 169. rufus, 257. violajugulum, 168, 396. Troglodytes aëdon, 24, 155, 164,

277. 326, 434. aëdon aztecus, 164, 400. aëdon marianæ, 163. aëdon parkmanii, 16, 163, 164, 362. alascensis, 410.

americana, 164. cahooni, 94. hiemalis, 153, 155, 388, 389,

hiemalis pacificus, 362. tobagensis, 404. Trogon ambiguus, 195.

Trumbull, G., notice of his 'Names and Portraits of Birds,' 414. Turdus aliciæ. 277, 310, 422.

aliciæ bicknelli, 153, 422. aonalaschkæ, 167, 365. aonalaschkæ auduboni, 167, 365.

aonalaschkæ pallasii. 153, 155, 277, 327, 328, 338, 389, 393.

celænops, 196. fuscescens, 151, 153, 155, 389. fuscescens salicicolus, 167. jouyi, 195. mustelinus, 25. ustulatus. 365.

swainsonii, 153, ustulatus

Turkey, Wild, 18, 204, 270, 314.

Turner, L. M., notice of his 'Con-

tributions to the Natural History of Alaska,' 409.

Turnstone, 110, 147. Turtur stimpsoni, 197.

Tympanuchus americanus. 349. cupido, 402. Tyrannus carolinensis, 200.

tyrannus, 21, 121, 149, 151, 154, 272, 389, 433. verticalis, 257.

ULULA, cinerea, 110. cinerea lapponica, 410.

Uria lomvia, 331. troile, 131-135, 281, 382.

Urinator imber, 140, 149, 268. 345, 373, 389, 390, lumme, 147, 170, 374.

VANELLUS chilensis, 290. Veery, 151, 153. Verdin, 166. Vireo alleni, 157. altiloquus barbatulus, 185, 186. belli pusillus, 33. caymanensis, 157. crassirostris flavescens, 157. flavifrons, 23, 275. flavoviridis, 210. gilvus, 23, 150, 152, 327, 389. gilvus swainsoni, 32, 258, 360, 399. huttoni stephensi, 32. noveboracensis, 23, 275. noveboracensis maynardi, 187, 399. olivaceus, 21, 23, 150, 152, 154, 275, 327, 389, 434-philadelphicus, 115, 152. solitarius, 152, 275, 389. solitarius alticola, 275, 324, solitarius cassinii, 32, 258, 360. solitarius plumbeus, 32. vicinior, 33. Vireo, Black-whiskered, 186. Blue-headed, 152. 275. Cassin's, 32, 258. Gray, 33. Key West, 187, 399.

Blue-headed, 152, 275.
Cassin's, 32, 258.
Gray, 33.
Key West, 187, 399.
Least, 33.
Mounta'n Solitary, 275, 399.
Philadelphia, 115, 152.
Plumbeous, 32.
Red-eyed, 23, 150, 152, 275.
Stephens's, 32.
Warbling, 23, 150, 152, 258.
Western Warbling, 32, 399.
White-eyed, 23, 275.
Yellow-throated, 23, 275.
Voorhees, C. G., a third specimen of Lawrence's Warbler, 427.

Vulture, Black, 270.

Turkey, 18, 179, 270.

WAGTAIL, Swinhoe's 410.

Warbler, Audubon's 34, 361.

Bachman's 187, 323, 325, 428.

Bay-breasted, 153, 433.

Black-and-white, 23, 152, 275.

Black-and-yellow, 153, 388,

Blackburnian, 150, 153, 433. Blackpoll, 153, 182, 276, 432. Warbler, Black-throated Blue, 149, 150, 153, 276, 391, 433. Black-throated Gray, 34, 104. Black-throated Green, 150, Blue, 431. Blue Yellow-backed, 150, 152. Blue-winged, 24. Calaveras, 33, 361. Canadian, 151, 153, 392, 393. Cape May, 187, 210, 275. Cerulean, 324, 431. Chestnut-sided, 150, 153, 433. Connecticut, 187, 323. Grace's, 34. Green Blackcap, 323. Hermit, 35 Hooded, 116, 276. Kentucky, 24. Kirtland's, 148. Lawrence's, 427. Lucy's, 33. Lutescent, 34. Macgillivray's, 35. Mourning, 153, 323. Myrtle, 276. Nashville, 150, 152, 323. Olive, 34. Orange-crowned, 275. Palm, 170, 182, 276. Parula, 24, 275, 431, 433. Pileolated, 36. Pine, 24, 276. Prairie, 24, 276, 405. Prothonotary, 170, 182, 210, 275, 322, 325. Red-faced, 36, 385. Sonora Yellow, 137, 400. Swainson's 187, 323, 325. Tennessee, 152, 187. Townsend's, 35. Virginia's, 33. Western Yellow, 34, 399. Worm-eating, 23, 275. Yellow, 16, 24, 152. Yellow Palm, 276. Yellow-rumped, 16, 34, 149, 150, 153, 388, 433.

Yellow-throated, 276. Water-thrush, 150, 153. Grinnell's, 35, 148, 324. Louisiana, 24, 276. Waxwing, Cedar, 23, 31, 275.

Waxwing, Cedar, 23, 31, 275. Wayne, A. T., habits of the Purple Gallinule (Ionornis martinica), 109; Loxia curvirostra minor again at Yemassee, S. C., 115; Loxia curvirostra minor taken again at Yemassee, S. C., 208; Ammodramus lecontei at Yemassee, S. C., 210; Ammodramus henslowii wintering in large numbers at Yemassee, S. C., 210.

Whip-poor-will, 21, 149, 151, 272. Eared, 89. Willet, Western, 185, 395.

Williams, R. S., Upper Missouri River birds, 14; spotted eggs of Parus gambeli, 118.

Wood, A. H., Ulula cinerea in Steuben Co., New York, 110. Woodcock, 20, 330, 331.

American, 270, 348. Woodpecker, Arctic Three-toed, 151. Downy, 16, 21, 151, 271.

Downy, 16, 21, 151, 271. Golden-winged, 149, 151. Hairy, 21, 149, 151. Harris's, 213, 214, 218, 252. Lewis's, 319. Pileated, 111, 149, 271, 313. Red-bellied, 21, 271. Red-cockaded, 271.

Red-headed, 21, 271. Red-naped, 216. Southern Hairy, 271. White-headed, 252

White-headed, 253. Yellow-bellied, 151, 325. Wren, Alaskan, 410.

Baird's, 163. Cactus, 116, 162. Cahoon's, 94. Cañon, 162. Wren, Carolina, 24, 183, 277.
Florida, 187.
House, 24, 277.
Long-billed Marsh, 24, 165.
Marian's, 163.
Marian's Marsh, 188, 400.
Parkman's, 163.
Rock, 162.
Tule, 400.

Western House, 16, 400. Winter, 153, 388, 392. Wood, 165.

XANTHOCEPHALUS xanthocephalus, 112, 262, 362, 414.

Xema atricilla, 75. furcata, 401. Xenopicus albolarvatus, 240, 253.

YELLOWBIRD, Summer, 150. Yellow-legs, 270, 349. Greater, 110, 349.

Yellow-throat, Maryland, 24, 151, 153, 276. Western, 35. Yunx torquilla, 212.

ZENAIDURA macroura, 17, 20, 119, 145, 270, 349, 389. Zonotrichia albicollis, 17, 149, 150,

> 426, 434. boucardi, 41. coronata, 358, 438-444. intermedia, 17, 358. leucophrys, 139, 181, 434. strigiceps, 412.

whitii, 412.

152, 154, 274, 388, 389, 390,

ERRATA.

Page 299, line 13 from top, for Metopidius indicus read Jacana indica. Page 346, lines 7 and 14 from top, for Mergus read Merganser.

